



Soil Degradation: Causes, Significant Threats and Management

Asha^{1*}, Diksha¹, Saloni
Yadav¹ and Sandeep Dagar²

¹Department of Soil Science,
CCS Haryana Agricultural
University- Hisar (HR.)

²Department of Vegetable
Science, CCS Haryana
Agricultural University - Hisar
(HR.)



Open Access

*Corresponding Author

Asha*

Article History

Received: 3. 01.2023

Revised: 8. 01.2023

Accepted: 12. 01.2023

This article is published under the
terms of the [Creative Commons
Attribution License 4.0.](https://creativecommons.org/licenses/by/4.0/)

INTRODUCTION

The term "soil degradation" describes a loss in the productivity of the soil brought on by unfavourable changes in the nutritional status, soil organic matter, structural characteristics, and concentrations of harmful compounds and electrolytes. The process of soil deterioration reduces the land's ability to provide commodities and services in the present as well as the future. It has also been described as the pace of unfavourable change in soil characteristics leading to a loss in the land's capacity for production because of processes primarily caused by human interference. As a result, it suggests a loss in soil productivity, a decline in plant cover, a qualitative and quantitative decline in water resources, soil degradation, and air pollution.

Degradation is a stage of evolution that lowers the potential of resources. 7.40 million hectares of arable land. Every year, due to climate change and deforestation, people around the world migrate to degraded regions. Since the beginning of land farming, soil degradation has been an issue. However, it has become significantly worse in recent decades as a result of India's population growth, which is expanding at a pace of roughly 1.8% and necessitating the plow-up of marginal regions to fulfil the rising food demand. These marginal soils are especially susceptible to deterioration, which lowers their quality and general production. In order to prevent soil degradation in India, the current paper briefly describes degraded and wastelands, their origins, significant threats, and management solutions.

Why Soil Degradation Occurs?

The overuse of the land to supply conflicting demands for food, fibre, and fodder from the expanding population is what is causing the destruction of the environment. Salinization, flooding, drought, erosion, and waterlogging are only a few of the human activities that exacerbate soil deterioration.

Other activities include the construction of large-scale irrigation canals, deforestation, and improper land use. In turn, these processes lower agricultural yield, which causes social insecurity.

The main cause of soil deterioration may be greenhouse gas emissions into the atmosphere, which cause global warming. Deforestation and removal of natural vegetation, excessive grazing, agriculture-related activities, and excessive domestic use of the vegetation are the causes of degradation brought on by direct and indirect human interference.

Major Threats and Management Options

Soil Erosion:- Soil erosion is a significant contributor to the depletion of the environment's natural resources. Soil erosion is typically more severe than in plain and undulating places where there are mountains. Improper soil management, unsuitable for Accelerated soil erosion is caused by things like tilling along a slope, not covering crops during heavy rains, etc. Land productivity is lost because of erosion.

Salinization and Alkalinization:- One of the primary tactics for reaching self-sufficiency in food production has been the extension of irrigation. From roughly 22 M hectares in 1950 to more than 51 M ha today, India's net irrigated area has risen. The majority of the development involves increasing the area irrigated by canals, which raises the groundwater table and causes the soil to degrade due to salt build-up. However, the area of 6 million ha of salinized or alkalinized soils has decreased due to reclamation operations from 7 million ha. Crop yields and output are affected by the high amounts of soluble salts, exchangeable sodium, or both present in these soils. The physiochemical characteristics and salt composition of the soils determine whether it is saline, sodic, or saline-sodic. The problem of salinity and sodicity in India will inevitably get worse if these waters are continuously used to irrigate agricultural crops.

Acidity:- The laterites and different latosolic soils, such as ferruginous red soils, ferruginous gravelly red soils, mixed red and black, or red and yellow soils, make up the greatest areas in India covered by acid soils. These soils developed because of extensive cation leaching brought on by heavy rainfall, which lowered the pH and reduced soil fertility. The classification of acidic soils as extremely acidic (pH 4.5), moderately acidic (pH 4.5-5.5), mildly acidic (pH 5.5-6.5), and non-acidic (pH > 6.5) soils is based on the pH range.


Acid soils present the normal water management issues that are mostly related to the physical and chemical characteristics of the soil. Acidic light-textured soils with a kaolinite dominance have extremely high-saturated hydraulic conductivity, which causes significant percolation losses.

Soil Organic Carbon Losses:- Due to nutrient depletion brought on by pedogenic processes for soil growth, Alfisols, Ultisols, and Oxisols are vulnerable to chemical degradation. These soils' base saturation is significantly lower than that of Alfisols, especially in Ultisols and Oxisols. The main causes of the depletion of soil organic carbon include intense farming, removal or in-situ burning of crop leftovers and minimal or no input of organic manures. To sequester organic carbon in soils, ideal choices include managing agricultural wastes, using inorganic and organic materials in a balanced and integrated manner, etc.

Nutrient Imbalance:- High agricultural yields require a balanced nutrient supply, yet too much or too little of a given nutrient can have negative effects on ecosystems, human health, and the environment. There are several different ways that nutrients can be lost, including through runoff, leaching, erosion, and emissions of NH₃, N₂O, NO, and N₂ into the air and discharges into water. Between Indian states and regions, nutrient inputs differ significantly. In agriculturally developed areas like Punjab, Haryana, and Orissa after the 1980s, fertiliser was more commonly used

than manure. The extensive use of fertiliser and the rapidly expanding animal industry have both contributed to significant N losses to the environment. Both the N inputs and outputs demonstrated. Large differences amongst crops were also seen in the N inputs and outputs without sacrificing crop yield, it is

possible to significantly reduce both the N imbalance and the N losses through the balanced and integrated use of organic manures and fertilisers, which are successful at boosting crop yields, improving nutrient use efficiency, and reducing environmental effects.



Socio-Economic Impacts and Future Challenges:-

The effects of soil deterioration on society and the environment are extensive. With over three-quarters of the poor and food insecure living in rural areas and depending on the natural resources for employment and income, it is now widely acknowledged that poverty in South Asia has historically been primarily a rural phenomenon. About 520 million people (40 per cent of the world's poor) live in South Asia, which has a poverty incidence of 43 per cent. Due to nutrient depletion caused by soil

erosion and other processes of soil degradation, the quality of the soil has decreased. People are forced to look for more land to farm because of poverty and the destruction of the environment. As a result, people are compelled to farm in woods, cultivate steep slopes, and further deteriorate marginal lands. Therefore, preserving natural resources and revitalizing wastelands and degraded areas present significant opportunities for reducing poverty and providing sustainable means of subsistence.