



The Plant's Secret Weapon: How GABA and BABA "Prime" Crops to Survive in Drought and Salt

Vikram Singh Gaur¹,
Anshuman Singh^{2*},
Chandu Singh³,
Rakesh Kumar⁴,
Chander Mohan⁵,
Ashutosh Sharma⁴

¹College of Agriculture, Balaghat, JNKVV, Jabalpur, M.P., India
²ICAR-IASRI, New Delhi
³ICAR-IARI, New Delhi
⁴RLB Central Agricultural University, Jhansi, U.P., India
⁵ICAR-IIMR, Regional Maize Research and SPC, Begusarai, Bihar, India



*Corresponding Author
Anshuman Singh*

Article History

Received: 1.12.2025

Revised: 5.12.2025

Accepted: 10.12.2025

This article is published under the terms of the [Creative Commons Attribution License 4.0](https://creativecommons.org/licenses/by/4.0/).

INTRODUCTION

The escalating effects of climate change, characterized by erratic rainfall and increasing soil salinization, pose a severe threat to food security worldwide. Traditional breeding methods to develop highly stress-tolerant crops are often slow and cannot keep pace with the rapidly changing environmental landscape. Consequently, there is a critical need for innovative and resource-efficient strategies to boost the natural resilience of staple crops. A fascinating concept emerging from plant biology research is "priming." Priming describes a state where a plant, having received a mild or transient stimulating signal (the priming agent), develops an enhanced capacity to mount a faster and stronger defense response upon subsequent exposure to a severe stressor. Crucially, the plant maintains this readiness without expending the significant energy required for a full defense response during non-stress conditions. Among the most effective and widely studied chemical priming agents are the non-protein amino acids, gamma-aminobutyric acid (GABA) and its isomer, beta-aminobutyric acid (BABA). While GABA is a ubiquitous, naturally accumulating metabolite found in all organisms and often linked to the plant's carbon/nitrogen balance, both compounds have been unequivocally identified as key signaling molecules that orchestrate the plant's stress "memory" and defensive readiness. Their application offers a non-toxic, eco-friendly approach to arming crops against the most challenging abiotic stresses: drought and salt.

2. The Dual Role of GABA in Stress Defense

GABA is synthesized in plants primarily via the GABA shunt, a metabolic pathway that bypasses parts of the conventional tricarboxylic acid (TCA) cycle. While this pathway is essential for general metabolism and acts as a carbon and nitrogen reservoir, GABA's role extends far beyond a simple metabolite, particularly under stress. Upon application or stress-induced accumulation, GABA acts as a crucial signal. It enhances stress tolerance through multiple mechanisms, including:

2.1. Antioxidant System Enhancement:

Drought and salinity typically lead to a surge in reactive oxygen species (ROS), causing oxidative damage. GABA priming accelerates the activation of key antioxidant enzymes, such as Superoxide Dismutase (SOD), Catalase (CAT), and Peroxidase (POD), effectively scavenging ROS and limiting cellular damage.

2.2. Osmotic Adjustment: Under drought and high salinity, plants must maintain cellular water pressure (turgor). GABA contributes to this by regulating the accumulation of compatible solutes, or osmolytes, like proline and soluble

sugars. Furthermore, GABA itself can function as a non-toxic osmolyte, helping to maintain cellular water balance and protect photosynthetic machinery.

2.3. Ion Homeostasis: In saline environments, the accumulation of toxic ions, particularly Na^+ , disrupts essential processes. GABA has been shown to modulate ion transport, aiding in the compartmentation of Na^+ ions away from the cytosol or promoting the uptake of beneficial ions like K^+ , thereby maintaining the critical K^+/Na^+ ratio necessary for cell function.




Figure-1: GABA role in alleviating salt stress negative effects

3. BABA: The Master Priming Agent for Resilience

BABA, a structural isomer of GABA, is known as one of the most potent chemical inducers of resistance, particularly for priming plants against both biotic (pathogens) and abiotic stresses. While it was once considered a foreign compound, it is now known to accumulate endogenously in stressed plants, confirming its role as a natural stress metabolite.

The priming effect of BABA is mediated through a complex signaling network. Research suggests that a specific aspartyl-tRNAsynthetase protein, encoded by the IBI1 gene, acts as the receptor for BABA. This receptor-ligand interaction triggers signaling cascades that crosstalk with major plant defense hormones:

3.1. Abscisic Acid (ABA): For abiotic stress, BABA priming is often linked to the ABA pathway, a central regulator of drought response. The priming action leads to a faster and more robust induction of ABA-responsive genes, which control critical drought-response mechanisms like stomatal closure and osmolyte synthesis.

3.2. Faster, Stronger Defense: Like GABA, BABA priming establishes an enhanced "stress memory." When a primed plant encounters drought or salt stress, its defense-related genes, including those for antioxidant production and ion transporters, are activated earlier and reach higher expression levels than in non-primed plants. This head-start minimizes the initial shock and damage caused by the stressor.

4. The Cost-Benefit Advantage of Priming

The most significant agricultural advantage of GABA and BABA priming is the cost-effective enhancement of tolerance. In the primed state, the plant is merely "ready," not actively fighting. Only a small amount of energy is invested in pre-synthesizing signaling components or modifying chromatin (epigenetic memory). The bulk of the energy-consuming defense machinery (like mass-producing antioxidants or defense proteins) is only mobilized rapidly upon the actual stress event. This conservation of metabolic resources ensures that the plant does not suffer growth penalties, which are common when traditional defense mechanisms are constitutively activated.

Plant Species	Abiotic stresses	Reference
Pistachio	Salt stress	Karimi, <i>et al.</i> , 2017
Rapeseed	Drought stress	Baghizadeh, <i>et al.</i> , 2017
Tobacco	Chilling stress	Ma, <i>et al.</i> , 2020
Chickpea	Salt stress	Mahmud, <i>et al.</i> , 2020
Chinese cabbage	Salt stress	Elradi, <i>et al.</i> , 2022
Linen, Chickpea	Heat stress drought stress	Quan, <i>et al.</i> , 2022

Table-1: Demonstration of many plant species that have been examined to elucidate the function of BABA in tolerance and regulation of different types of stresses.

CONCLUSION

GABA and BABA represent a paradigm shift in crop protection against environmental extremes. Their function as signaling molecules that induce an energy-efficient state of stress priming offers a powerful solution to mitigating the damaging effects of drought and salinity. By accelerating the activation of core defensive systems—from antioxidant scavenging and better osmoregulation to maintaining crucial ion balance—these non-protein amino acids allow crops to maintain higher yield and survival rates under adverse conditions. Future research focusing on optimizing the application methods and better understanding the receptor-level signaling of these molecules promises to unlock their full potential, establishing them as the plant's essential "secret weapon" for a more sustainable and resilient agriculture.

REFERENCES

Baghizadeh, A. (2017). Alleviation of oxidative stress induced by drought stress through priming by β -aminobutyric acid (BABA) in Rapeseed (*Brassica napus* L.) plants. *Plant Physiology*.

Bouche, N. and Fromm, H. (2004). GABA in plants: Just a metabolite? *Trends in Plant Science*, 9(3), 110–115.

Decsi, K., Ahmed, M., Rizk, R., Abdul-Hamid, D., Kovacs, G. P. and Toth, Z. (2024). Emerging trends in non-protein amino acids as potential priming agents: implications for stress management strategies and unveiling their regulatory functions. *International Journal of Molecular Sciences*, 25(11), 6203.

Elradi, S., Suliman, M., Zhou, G., Nimir, E., Nimir, N., Zhu, G. and Ali, A. (2022). Seeds priming with β -aminobutyric acid alleviated salinity stress of chickpea at germination and early seedling growth. *Chilean journal of agricultural research*, 82(3), 426-436.

Jakab, G., Manz, G., Mauch-Mani, B., Baltruschat, H., Kogel, K.H., Krogge, W. and Mauch, F. (2005). Beta-Aminobutyric acid-induced resistance in *Arabidopsis thaliana* in the absence of priming for pathogenesis-related gene expression and salicylic acid accumulation. *European Journal of Plant Pathology*, 112(1), 1–11.

Jisha, K.C., Shackira, A.M., and Puthur, J.T. (2018). GABA/BABA Priming Causes Signaling of Defense Pathways Related to Abiotic Stress Tolerance in Plants. *Neurotransmitters in Plants*, CRC Press.

Karimi, S., Mirfattahi, Z., Ferguson, L., & Tavallali, V. (2017). Using controlled salt stress and β -aminobutyric acid signaling to decrease transplant failure. *ScientiaHorticulturae*, 225, 156–162.

Luna, E., Pastor, V., Robert, J., Flors, V., Mauch-Mani, B. and Ton, J. (2014). Callose deposition: the first line of defense in the BABA-induced priming of *Arabidopsis*. *Molecular Plant-Microbe Interactions*, 27(12), 1403–1413.

Ma, Y., Dias, M. C., & Freitas, H. (2020). Drought and salinity stress responses and microbe-induced tolerance in plants. *Frontiers in plant science*, 11, 591911.

Mahmud, J. A., Hasanuzzaman, M., Khan, M. I. R., Nahar, K. and Fujita, M. (2020). β -Aminobutyric acid pretreatment confers

salt stress tolerance in *Brassica napus* L. by modulating reactive oxygen species metabolism and methylglyoxal detoxification. *Plants*, 9(2), 241.

Quan, J., Zheng, W., Wu, M., Shen, Z., Tan, J., Li, Z. and Zang, Y. (2022). Glycine betaine and β -aminobutyric acid mitigate the detrimental effects of heat stress on Chinese cabbage (*Brassica rapa* L. ssp. *pekinensis*) seedlings with improved photosynthetic performance and antioxidant system. *Plants*, 11(9), 1213.

Ramesh, S.A., Tyerman, S.D., Gillham, M. and Xu, B. (2017). Gamma-aminobutyric acid (GABA) signaling in plants. *Cellular and Molecular Life Sciences*, 74(9), 1577–1603.

Wang, J.Y., Zhang, Y.K., Li, Q., Zong, J.S. and Lu, Z.Y. (2017). Exogenous gamma-aminobutyric acid (GABA) enhances the salt tolerance of maize by improving photosynthesis, osmotic adjustment, and antioxidant defense. *Photosynthetica*, 55(1), 164–173.

Yasir, T., Ateeq, M., Wasaya, A., Hussain, M., Sarwar, N., Mubeen, K. and Sabagh, A. (2023). Seed Priming and Foliar Supplementation with β -aminobutyric Acid Alleviates Drought Stress through Mitigation of Oxidative Stress and Enhancement of Antioxidant Defense in Linseed. *Phyton*, 92(11), 3113.