

Climate Change and Its Impact on Insect Pests

Priti Priya^{*1},
Manoj Baranwal²

¹Assistant Professor Cum Junior
Scientist, Birsa Agriculture
University, Ranchi Pin
Code-834006

²Sr. Scientist-Cum-Associate
Professor, Plant Pathology,
BAU, Ranchi- 834006



Open Access

*Corresponding Author
Priti Priya*

Article History

Received: 2. 4. 2026

Revised: 6. 4. 2026

Accepted: 11. 4. 2026

This article is published under the
terms of the [Creative Commons
Attribution License 4.0](https://creativecommons.org/licenses/by/4.0/).

INTRODUCTION

Climate change refers to long-term and significant alterations in climatic variables such as temperature, rainfall, humidity, and atmospheric composition. Human activities which include greenhouse gas emissions and deforestation and industrialization drive these environmental changes. The agricultural sector depends on climate conditions because they control every aspect of crop growth from development through productivity.

Crop production faces many biotic stresses which include insect pests that demonstrate particular susceptibility to climate change because they possess high reproductive capacity and quick life cycle and their environmental reactions show strong sensitivity. Insect pests cause substantial yield losses in both field crops and horticultural crops which affects farmers because it brings economic losses. The changing climate patterns are causing pests to change their behavior and population patterns and distribution. Climate change will create more severe pest problems which will make pest control activities more difficult and complex.



2. Key Climatic Factors Affecting Insect Pests

2.1 Temperature

Temperature serves as the main climatic element which determines how insects grow and develop and manage their life span. Most insects are classified as poikilothermic (cold-blooded) because their body temperature and bodily functions depend on the surrounding temperature.

The body requires more energy when temperatures rise because the heat speeds up all chemical reactions in the body. The phenomenon of multivoltinism occurs when insects which can produce more than one generation during a single year. The increase in generations leads to greater population growth which results in more pest pressure against crops. Insects expand their geographic territory because rising temperatures permit them to enter new areas which they could not reach before because of cooler climatic conditions.

2.2 Carbon Dioxide (CO₂) Concentration

The rising atmospheric CO₂ levels create indirect but major impacts on insect pests because they change how plants grow and their nutritional content. The presence of increased CO₂ gas leads to enhanced plant growth which results in higher biomass production but this development process decreases the nitrogen levels found in plant tissues.

The decreasing nutritional content of plants requires herbivorous insects to consume larger amounts of plant material because they need to obtain their necessary nutrients. The increased feeding behavior of pests leads to more severe agricultural damage which affects crop production. The chemical composition of plants undergoes changes which affect their ability to defend themselves thus making their crops more vulnerable to pest attacks.

2.3 Rainfall and Humidity

Insect populations depend on two essential factors which are rainfall and humidity. Insects that have soft bodies which include aphids and mites and whiteflies find high humidity conditions to be optimal for their growth and reproduction.

The process of heavy rainfall creates two main impacts because it removes insect populations through physical washing and it diminishes their overall population size. The combination of irregular rainfall patterns and

extended periods of drought followed by intense rainfall creates conditions which enable pests to multiply. The natural equilibrium experiences interruption because of these environmental changes which result in quick pest population growth after the destructive conditions end.

2.4 Extreme Weather Events

The climate change crisis has increased the frequency of extreme weather events which now include droughts and floods and cyclones and heatwaves. These incidents create significant impacts which influence both insect pests and their natural enemies.

Extreme weather conditions decrease pest numbers for a short time interval but they create ecological disturbances because they harm predator and parasitoid populations more than pest species. The pest population will experience a resurgence after the conditions have turned beneficial again. The condition of drought activates stress on plants which results in decreased plant strength and increased vulnerability to insect infestations.

3. Impact of Climate Change

Impact of Climate Change on Insect Pest Dynamics contains two parts which explain how climate change has affected insect pest distribution patterns throughout the world.

3.1 Changes in Geographical Distribution

presents evidence that climate change creates current geographical. Rising temperatures enable pests to travel beyond their previous survival limits towards higher altitudes and higher latitudes.

The new pest expansion brings fresh challenges to areas which have never experienced certain insect species before. Current agricultural practices need to adapt because tropical and subtropical pests have begun to spread their range into temperate regions.

3.2 Increased Pest Survival and Overwintering

Warmer winters reduce the mortality of insect pests that typically die under extreme cold conditions. The population of insects which survive off-season periods will enter the new growing season with their peak numbers.

The crop damage risk increases because early-season infestations develop into more severe problems. Pests build rapid population growth during the growing season because of their success in surviving the winter months.

3.3 Altered Life Cycle and Phenology

Climate change affects the timing of biological events, known as phenology, in insect pests. Organisms develop faster when temperatures increase which leads to them emerging earlier and producing shorter life cycles.

Pests establish better life cycle synchronization with their host plants which results in improved feeding efficiency and increased damage. The synchronization permits pests to attack crops when the plants experience their most critical vulnerabilities.

3.4 Pest Outbreaks and Resurgence

Pest outbreaks occur suddenly when climate conditions differ from their normal patterns and extreme weather events happen. Environmental conditions that change frequently create disturbances which prevent predators and parasitoids from executing their natural pest control functions.

Pest populations experience rapid growth because biological control systems fail to function effectively after their initial control period ends. Uncontrolled outbreaks result in severe crop damage which requires effective management to prevent.

3.5 Emergence of New Pest Species

Climate change enables invasive alien species to invade and establish themselves in new ecosystems. Invasive pests gain a competitive edge because they enter environments which have not developed defenses against their presence.

The emergence of new pest species creates additional challenges for farmers and researchers because current management practices fail to handle these unidentified pests.

4. Impact on Crop Production

The changing dynamics of insect pests have direct and indirect effects on crop production. Increased pest pressure leads to significant yield losses and deterioration in crop quality.

Farmers increase their dependence on chemical pesticides because they need to handle pest problems which results in higher expense and environmental danger. Pesticides need to be used at limited amounts because their excessive application results in two problems. Pesticide overuse creates two major problems which threaten food production systems and farmers' ability to earn a living.

5. Impact on Natural Enemies

The essential function of natural enemies through their predation and parasitoid

activities and their pathogen attacks leads to effective insect pest population control. But climate change brings negative effects to these beneficial organisms.

The different ways pests and their natural enemies respond to climate changes create development patterns which take place at different times thus decreasing biological control efficiency. The decline of natural enemy populations occurs when biodiversity and habitat conditions change which results in decreased ecosystem stability.

6. Case Examples

Real-world situations demonstrate how climate change affects insect pest populations. The *Helicoverpa armigera* (cotton bollworm) population has increased in warmer climates because those conditions enable its growth and reproduction.

Whiteflies and aphids have spread to new areas while becoming major pest threats in vegetable crops. Under the current climatic changes in rice ecosystems stem borers show greater pest activity which leads to heavier crop damage.

7. Adaptive Strategies for Pest Management

7.1 Integrated Pest Management (IPM)

Sustainable pest management requires the implementation of Integrated Pest Management (IPM) strategies. Through its combination of biological and cultural and mechanical and chemical methods IPM system protects against pests while decreasing environmental damage. The system gains strength through the combination of pest-resistant varieties and ecological methods.

7.2 Climate-Smart Agriculture

Climate-smart agricultural practices involve adjusting cropping patterns and sowing dates with crop selection to decrease pest threats. The combination of crop diversification and intercropping systems leads to decreased pest problems and enhanced system stability.

7.3 Monitoring and Forecasting

Pest population monitoring needs to be done continuously through regular surveillance because it enables quick response when intervention is necessary. Farmers use weather-based pest forecasting models and early warning systems to implement preventive actions that stop pest outbreaks from happening.

7.4 Biological Control Enhancement

Natural enemy conservation together with their population increase serves as the

foundation for effective pest control systems. Biological control practices see improvements through habitat management practices which decrease pesticide usage while biocontrol agents are released.

7.5 Use of Digital Technologies

Pest management systems undergo transformation through the application of remote sensing technology together with artificial intelligence and mobile advisory systems. The systems provide farmers with capabilities to monitor pest activity in real-time while they can precisely forecast pest behavior and receive urgent alerts.

8. Future Research Needs

Research should develop pest management techniques which can withstand climate variations and environmental changes. Scientists need to study how climate-related stressors affect pest species evolution and their adaptive capacity to identify their climate adaptation techniques. Scientists need to improve pest forecasting models through the integration of climatic data and biological information together with ecological research findings. Sustainable pest control methods which protect the environment will play an essential role in achieving sustainable agricultural practices.

CONCLUSION

Insect pest behavior patterns undergo fundamental transformations because of climate change which creates major obstacles for agricultural productivity and sustainability efforts. The agricultural industry needs to create active pest management systems which can handle the increasing number and strength of pest outbreaks.

Effective insect pest management under climate change requires the integration of traditional knowledge together with contemporary technologies and adaptive strategies which can withstand future climate changes.

REFERENCES

- Kambrekar, D. N., Guledgudda, S. S., Katti, A., & Kumar, M. (2015). Impact of climate change on insect pests and their natural enemies. *Karnataka Journal of Agricultural Sciences*, 28(5), 814-816.
- Karuppaiah, V., & Sujayanad, G. K. (2012). Impact of climate change on population dynamics of insect pests. *World Journal of Agricultural Sciences*, 8(3), 240-246.
- Laštůvka, Z. (2009). Climate change and its possible influence on the occurrence and importance of insect pests. *Plant Protection Science*, 45(Special Issue), S53-S62.
- Nayak, S. B., Rao, K. S., & Ramalakshmi, V. (2020). Impact of climate change on insect pests and their natural enemies. *Int. J. Ecol. Environ. Sci*, 2, 579-584.
- Pareek, A., Meena, B. M., Sharma, S., Tatarwal, M. L., Kalyan, R. K., & Meena, B. L. (2017). Impact of climate change on insect pests and their management strategies. *Climate change and sustainable agriculture*, 253-286.
- Skendžić, S., Zovko, M., Živković, I. P., Lešić, V., & Lemić, D. (2021). The impact of climate change on agricultural insect pests. *Insects*, 12(5), 440.