

Organic Farming and Insect Pest Management: Challenges and Opportunities

Shahin Khan¹, Kritika Saklani², R.K Mishra³, Niveditha M.P⁴, Ramnivas⁵

¹Scientist, Department of Entomology, RVSKVV, ZARS Khargone, Madhya Pradesh.

²Ph.D Scholar, Department of Entomology, Punjab Agricultural University, Ludhiana.

³Assistant Professor, Department of Forestry, Doon (PG) college of Agriculture Science and Technology Camp Road Selaqui, Dehradun, Uttarakhand.

⁴Assistant Professor, Department of Agronomy, ZAHRS Mudigere KSNUAHS, Shivamogga.

⁵Assistant Professor, Department of Ag. Entomology, Maharaja Surajmal Agriculture College Rahimpur, Bharatpur (Raj.).



*Corresponding Author
Shahin Khan*

Article History

Received: 31.05.2026

Revised: 05.06.2026

Accepted: 10.06.2026

This article is published under the terms of the [Creative Commons Attribution License 4.0](https://creativecommons.org/licenses/by/4.0/).

INTRODUCTION

Organic agriculture is a production system that is sustainable and promotes environmental stewardship, is based on whole-ecosystem management principles, relies on biodiversity for pest management and nutrient cycling, and makes minimal use of synthetic inputs, including fertilizers and pesticides. According to the Food and Agriculture Organization (FAO), organic farming depends on ecological processes that bolster biodiversity and locally adapted cycles, that does not rely on external sources of chemicals, representing a cornerstone in sustainable food systems.

Insect pest management continues to be a major challenge for organic production systems as synthetic insecticides are prohibited, but it is clearly one of the most important problems to solve. Consequently, farmers must rely on a combination of preventive, ecological, cultural, and biological approaches to manage pest populations and protect crop yields. Organic farming emphasizes the resilience of the whole ecosystem and its natural pest control methods, instead of conventional agriculture's focus, which is to achieve controlled pace of pests through chemical means.

Having a rapid growth with more and more worldwide demand for safe, residue free and environment friendly food products, the organic agriculture practice is now extending all over the world. This expansion has increased the demand for insect pest management approaches that are efficient, dependable, and economically sustainable to maintain high yields in organic production systems.




2. Principles of Organic Pest Management

Organic insect pest management is based on the principles of ecological balance and prevention rather than eradication.

2.1 Prevention

Utilization of resistant crop varieties, crop rotation and intercropping, appropriate sowing and harvesting time, and soil fertility management are the main preventive measures in sustainable crop production system. Use of resistant varieties contribute in reducing pest and diseases incidences by having genetic resistance, which leads to the less vulnerability of the crop and less use of chemicals. Rotation


and intercropping promote higher levels of agroecosystem diversity and disrupt pest life cycles by disconnecting host continuity, so pest populations are reduced. Timely sowing and harvesting enable crops to avoid the peak periods of pest damage, which not only reduces the damage to crops but also facilitates harvest in a stable production. Soil health Based on Information in the abstracts Improvement of soil health through the addition of organic materials, manure, compost, and bio-fertilizers stimulates soil microbial activity and nutrient availability, which may improve plant Vigor and ability to withstand attacks by pests.



2.2 Conservation of Natural Enemies

Biocontrol tactics include enhancement of natural enemies (e.g., ladybird beetles, lacewings), preservation of parasitoids (e.g., *Trichogramma spp.*), and habitat diversification. Natural enemies of pests, such as predators (which consume larvae and adults


of pests like aphids, and whiteflies) and parasitoids (which parasitize pest eggs) suppress pest populations. The habitat diversification also enabled beneficial insects to find forage and shelter, and enhanced natural biocontrol for an ecological equilibrium.



2.3 Monitoring and Thresholds

Regular field scouting, use of pheromone and sticky traps, and decision-making based on pest population levels are essential components of effective pest monitoring and management. Regular field scouting helps in early detection of pest incidence and assessment of crop health status. Pheromone and sticky traps assist in monitoring pest activity and population

dynamics, enabling timely intervention. Decision-making based on economic threshold levels ensures that control measures are applied only when pest populations reach damaging levels, thereby avoiding unnecessary interventions and promoting sustainable pest management.



2.4 Ecological Balance

Preservation of agroecosystem diversity and stimulation of plant-insect-environment interactions are fundamental to ecological regulation of pests. The diversity of the agroecosystem provides habitat for numerous organisms, including natural enemies of pests, enhancing biological control. Improved plant-insect-environment interactions Favor ecosystem stability, pest outbreaks and amplify natural regulatory processes in the agroecosystem.

3. Major Insect Pest Problems in Organic Farming

Insect pests in agriculture can be broadly categorized based on their feeding habits and damage patterns. Chewing pests such as caterpillars (*Helicoverpa armigera*), leaf beetles and armyworms damage crops by feeding on leaves, reducing photosynthetic area and overall plant vigor. Sap-sucking pests including aphids, whiteflies, thrips and jassids extract plant sap, leading to nutrient loss, stunted growth and transmission of viral diseases. Borers and internal feeders such as fruit borers in tomato and cotton and stem borers in cereals attack internal plant tissues, making them difficult to detect and control. Polyphagous pests like fall armyworm (*Spodoptera frugiperda*) and mealybugs have a wide host range and can cause severe and widespread damage. These pests collectively contribute to significant yield losses if not managed through integrated and timely pest management strategies.

4. Challenges in Organic Insect Pest Management

Organic insect pest control is limited by several important factors. A major constraint is the unavailability of rapidly effective curative measures similar to synthetic insecticides. Organic systems are also susceptible to pest outbreaks because of lack of early suppression tools, conducive microclimatic conditions for pest development and monoculture-based

organic culture practices. In addition, farmers knowledge and skill-levels in pest identification, use of biological control measures, and ecological monitoring limits their capacity to manage pest effectively. Adoption is also limited by economic factors such as increased labour demand, expense of biopesticides and traps, and by the unpredictability of yields during the first years of conversion. The activity of neem or other botanicals-based products varies and the supply of good quality bio-control agents is limited, which compounds the difficulty. In addition, climate change intensifies pest challenges by accelerating their reproduction rates, facilitating the introduction of invasive species and altering predator-prey interactions in agricultural systems.

5. Opportunities in Organic Pest Management

Organic farming systems provide several innovative and sustainable opportunities for effective insect pest management despite existing challenges. Biological control remains the backbone of organic pest management, involving the use of parasitoids such as *Trichogramma* and *Bracon* spp., predators like ladybird beetles and spiders, and microbial pathogens including *Bacillus thuringiensis* (Bt), *Beauveria bassiana*, and *Metarhizium anisopliae*, all of which contribute to long-term pest population suppression. Botanical pesticides derived from plants such as neem (*Azadirachta indica*), pongamia oil, pyrethrum, and extracts of garlic and chili are widely used, with neem formulations being particularly effective as antifeedants and growth regulators. Habitat management and agroecological practices such as flower strips, mixed cropping, intercropping, agroforestry systems and hedgerow conservation promote biodiversity and enhance natural pest control. Pheromone and trap techniques, e.g. sex pheromone traps, light traps and hanging sticky traps, are cost-effective and environmentally safe instruments to monitor and mitigate pest

densities. Integrated Organic Pest Management (IOPM), a synthesis of cultural, biological, mechanical and botanical approaches with emphasis on prevention and control at threshold, increases system robustness and decreases reliability on standalone control methods. Moreover, digital and precision agriculture solutions such as drone-based surveillance, AI-powered models for pest prediction, m- Advisory systems, and remote sensing tools are increasingly being used to facilitate early detection and timely interventions for pest management, which also contributes to reduced crop losses and enhanced productivity.

6. Future Prospects

The prospects for organic insect pest management have never been brighter with progress in agroecology and biotechnology. Notable future challenges will be to produce climate-resilient biocontrol agents that are effective under altered environmental conditions, and to develop more CRISPR-based pest-resistant crop cultivars that hold organic principles by minimizing use of external inputs. The mass production of entomopathogens (e.g., *Bt*, *Beauveria*, *Metarhizium*) in biofactories will further increase the access to and cost-effectiveness of

biological control methods. Policy incentives for organic inputs usage will also be required to support farmers in adoption. Also, farmer field schools and extension services need to be strengthened to raise awareness, skill and uptake of integrated pest management practices. By merging traditional ecological knowledge with scientific advancements, it will contribute significantly in establishing adaptive, sustainable, organic pest management strategies.

CONCLUSION

Organic farming provides a viable alternative to conventional agriculture, yet insect pest control is still a critical constraint. Although challenges, for example, pest outbreaks, curative options are limited and there are gaps in knowledge, new opportunities are developing related to biological control, botanical pesticides, habitat management and digital technologies. For long-term success, a systems-based IOPM approach focusing on ecological balance, prevention and conservation of biodiversity is needed. Robust institutional support, farmer capacity building and research innovation will be the key to global expansion of organic pest management practice.